





BÁBOLNA **TETRA-SL**
COMMERCIAL LAYER
MANAGEMENT GUIDE



**BÁBOLNA TETRA-SL
COMMERCIAL LAYER
MANAGEMENT GUIDE**

CONTENTS

Introduction	2
Performance Specifications	3
Husbandry	4
Temperature	4
Stocking Rate	5
Feeding Space	5
Drinking Space	5
Beak Trimming	6
Water Intake	7
Light Intensity	7
Nutrition and Body Weight Control	8
Body Weight Control	8
Feed Intake	10
Body Weight Guide	11
Feeding Program	12
Nutritional Recommendations	13
Lighting Program	14
Veterinary Control	16
Recommendation for Vitamin Treatments	16
Egg Numbers	17
Rate of Lay	18
Egg Weight	19
Egg Weight and Size	20
Egg Quality	21
Egg Production and Egg Weight Graph	22
Hot Climate Management	23



INTRODUCTION

This guide contains information, which will help achieve the potential performance from TETRA-SL stock. It is, however, only a guide. The actual performance obtained will depend on a whole list of factors. Health status, ambient temperature, management and housing system are just some of the influences, which will modify the genetic potential bred into the TETRA-SL.

The economics of egg production vary around the world and from one part of a year to another. The attainment of maximum egg numbers may not result in maximum profit if egg price differentials indicate that management procedures should have been designed to produce maximum egg weight. TETRA-SL is an extremely versatile brown egg-laying hybrid, and management procedures may be modified so that it produces the correct balance of egg numbers and egg weight to suit a particular set of economic conditions.

Under each production parameter there is a summary of the effects that some of the more important management factors have upon performance. We hope that these summaries, which include recent research findings, will give the necessary information to enable the management protocol most appropriate for maximising profit from your TETRA-SL to be chosen.

Laying hens are living creatures, and each is an individual, it is therefore not possible to give any guarantee of performance, even if the management procedures described in this guide are followed precisely. However, adherence to the various recommendations will ensure that the best possible performance under the circumstances, may be attained.

Maximum performance will only be obtained when birds are kept in controlled environment conditions. However, it is possible to adapt management, when birds are kept in tropical open housing, to minimize the loss of performance caused by high temperature and natural lighting conditions. This guide includes a section devoted to the management of TETRA-SL in hot climate open housing.

Bábolna TETRA Ltd.

PERFORMANCE SPECIFICATIONS (STANDARD MATURITY)

Data to 80 weeks of age in controlled environment conditions

Liveability

0 - 17 weeks of age	97-98%
17 - 80 weeks of age	94-96%

Feed Intake

0 - 17 weeks of age	5.8-6.0 kg
17- 80 weeks of age	110-115 g/day

Body Weight

At 17 weeks of age	1.44 kg
at 80 weeks of age	1.92-2.00 kg

Sexual Maturity

Age at 50% rate of lay	144 days
Age at 90% rate of lay	159 days

Egg Production

Peak production	95-96%
Egg production above 90%	16-20 weeks
Total egg production for 52 weeks of lay	319 eggs
Total egg production for 80 weeks of age	363 eggs

Egg Mass Output

Total egg production for 52 weeks of lay	20.5 kg
Total egg production for 80 weeks of age	23.5 kg
Average egg weight at 80 weeks of age	67.7 g

These specifications assume an average ambient house temperature of 20°C (68°F)

It is important to note that in some countries welfare regulations may stipulate stocking rates and feeding and drinking spaces, which are different to those given in this manual. Regulations may also prohibit or restrict certain husbandry practices.

Things to do before delivery of chicks

The risk of an infection of any kind of poultry disease can be minimized by isolating a flock from other especially older flocks and by avoiding mixed-aged-flocks on the same farm.

All the building interior, including the drinking, feeding, heating and ventilation systems as well as the cages or slats and also the attached service areas and equipment has to be cleaned, disinfected and dried properly.

After reinstalling the disinfected and dried equipments they have to be checked whether they work properly and are adjusted for the right height.

Traps or poison for mice and flies have to be placed inside the building out of reach of the birds. Windows have to be covered by nets to keep wild birds outside the building.

Once the farm is disinfected and ready for a new flock, the entrance of unauthorised people and vehicles has to be minimized.

24 hours before delivery the following things have to be done:


- Start heating to reach the required temperature
- Check the drinking system and water temperature
- Prepare feed ready for the birds
- Check appropriate light intensity

Temperature

During the first 3 or 4 days chickens need 32-34 C° depending on the brooding system. To reach the required

temperature heating has to be started 24 hours prior to placing the day-old flock. The behaviour of the chickens is the best indicator of the temperature especially during night. If the birds are calm and quiet and they spread out equally in the house it means they feel comfortable. As the chicks are growing the temperature can be reduced to 30-32 C° by the end of the first week. From the second week the temperature can be reduced by 2-3 C° weekly until reaching 20 C°. Always measure the temperature at the bird's level. Besides the temperature it is essential to maintain proper humidity as well especially if brooding in cages. Relative humidity has to be kept between 40-60% by evaporating water (floor brooding) or watering the walks (cage brooding) if necessary.

Brooding temperature		
Age	Floor Brooding	Cage Brooding
1-4	32-33 C°	33-34 C°
5-7	30 C°	32 C°
8-14	28 C°	30 C°
15-21	26 C°	27 C°
22-28	24 C°	24 C°
29-35	22 C°	22 C°
36-119	20 C°	20 C°



Behaviour of the chicks is the best indicator of the appropriate temperature

Stocking Rate

(a) Floor systems

Environmental factors, such as type of housing, ventilation and temperature, have a greater effect upon stocking rate than genetic make-up. Slatted floors, for example, will allow a tighter stocking rate than litter, while high temperature, especially if combined with high relative humidity, necessitates a more liberal stocking rate. The following recommendations are given as a guide for litter units with an average temperature of about 20°C (68°F) at bird height. These rates should be reduced by 2% for each 1°C (2°F) rise in temperature above 20°C (68°F).

Age (weeks)	Deep litter	Slats
	Birds per m ²	
0-8	15	16
9-18	10	12
>18	7	8

(b) Cage systems

Stocking rates in cage systems are mainly determined by body weight and temperature.

Age (weeks)	Birds per m ²	Area per bird (cm ²)
0-8	66	150
9-18	29	350
>18	18	550

Feeding Space

Insufficient feeding space during rearing will produce uneven birds at sexual maturity, and result in reduced egg output in lay. The following should be regarded as the minimum requirements for satisfactory performance.

(a) Floor systems

Age (weeks)	Feeder space per bird (cm)	Birds per 40 cm dia circular feeder
0-2	3.5	70
	plus extra feeders in first week	
3-18	7.0	35
>18	10.0	30

(b) Cage systems

Age (weeks)	Feeder space per bird (cm)
0-2	2.0
	plus extra feeders in first 3-5 days
3-8	4.0
9-18	8.0
>18	10.0

Drinking Space

Water is an essential nutrient by itself. It can also influence all other nutrient intakes by controlling feed intake. For example, a restriction on water intake will cause a voluntary reduction in feed intake. Therefore the provision of an adequate number of well distributed drinking points is a key factor in egg production. In hot climates the need for an adequate supply of drinkers is even more important, as at high temperatures evaporative cooling plays a dominant role in the maintenance of a normal body temperature.

The drinking space recommended below must be increased in hot climates, and when water control is practised. In these situations there will be abnormally heavy demands for water at certain times of the day. To ensure that all birds find water when initially housed there should be a minimum light intensity of 20 lux at bird level. This is especially important at day old and where a change of drinking system occurs when moving birds into the laying house. It is also



recommended that extra drinkers are provided during the first week of life to minimize the incidence of non-starters or starve-outs.

(a) Floor systems

Age (weeks)	Birds per nipple	Trough space per bird (cm)
0-2	8 plus extra fountains (50 birds per fountain)	fountains
3-18	8	2.5
>18	8	4.5

(b) Cage systems

It is recommended that nipple and cup drinkers are located so that each cage of birds has access to at least 2 drinkers.

Age (weeks)	Birds per nipple	Trough space per bird (cm)
0-8	15	2.5
9-18	8	5.0
>18	4	10.0

Beak Trimming

Beak trimming (often incorrectly termed debeaking) need not be carried out routinely when TETRA-SL birds are kept in controlled environment housing. But if experience from previous flocks suggests that it is necessary, it will be worthwhile first checking all other aspects of management, before embarking on a program of beak trimming. The provision of more feeders and drinkers, more space per bird or improved ventilation may be the correct action. However, in open sided housing, routine beak trimming is recommended, as both bright light intensities and high temperatures may predispose undesirable behaviour. When essential,

beak trimming is best performed with a precision machine between 6 and 8 days of age. Only healthy birds can be beak trimmed. Electrolytes and vitamin K should be used in the drinking water before and after the procedure as well as deeper feed in the feeders.

Beak trimming, when done correctly, minimizes feed wastage and checks undesirable behaviour. It does not directly affect egg production. However, if performed incorrectly, it will reduce feed intake and, in turn, adversely affect egg output. Faulty techniques include trimming at the wrong age, cutting only one mandible, the removal of too much beak and unsatisfactory cauterization.

Care must be taken that all birds are correctly and uniformly beak trimmed. Each bird should mature with a rounded, but slightly shortened beak and well able to conduct normal feeding activity.



It is also important that the operation is conducted in compliance with any welfare regulations, which may exist. In some countries the procedure may be prohibited.

Water intake at 20 C°

Age (weeks)	Daily water intake (litre per 100 birds)
1	2
2	4
4	6
6	8
8	10
10	12
12	14
14	16
16	18
18	20
20	22
22	25

Light Intensity

Genetic progress, as well as changes in ration specifications, appear to have contributed to the ability of the modern egg-laying hen to perform perfectly satisfactorily at light intensities that would have been considered too low in the past. This gives the poultry farmer the opportunity of reducing light intensity, rather than beak trimming, to control undesirable bird behaviour.

Management, and in some countries welfare regulations, require the light intensity to be sufficiently bright to properly inspect the flock for health and remove dead and sick birds. The light intensity should also be high enough to check that the drinking system is functioning correctly and that the house is acceptably clean.

A light intensity of 10 lux (1 foot candle) in the laying house at feed trough level seems to be a satisfactory compromise between the need to control bird behaviour and feed wastage, and the requirements for bird inspection and maximization of egg output.

Chick Starter (0 - 8 weeks)

Essentially a Starter ration aims to produce a good skeleton, good organ development and help promote an active immune system. This is achieved by feeding the starter ad libitum with the correct balance and absolute levels of essential amino acids, and a minimum concentration (1.25%) of linoleic acid for growth, development of the immune system, feathering and skin condition.

Normally it is adequate to feed the Starter ration for 6 weeks, however, if for whatever reason body weight is substantially less than 500 grams at 6 weeks it is advisable to continue feeding Starter until body weight is on target.


Grower (8 - 15)

Whilst the Grower ration will be the lowest density ration that the bird receives, it is important that all nutrients are correctly included. Feed restriction is not recommended during this period as it may be difficult to achieve the correct body weight at first egg, particularly if early sexual maturity is desired. Feeding ad libitum is suggested but it is essential to monitor the body weights weekly during this period.

Body weight control

Body weight, especially prior to first egg, is a very important factor in determining the performance that will be achieved during the laying period. Therefore a regular body weight control is essential. At least 2% of a flock has to be measured not less than every two weeks. Birds have to be weighed in the morning hours and on the same day of the week before feeding.

The growth of a flock is normal and the birds can be considered equal if the difference between the individual and average weight is not more than 10%.



At least 2% of a flock has to be measured

As TETRA-SL pullets are not liable to overweight, ad libitum feeding can be used during the whole rearing period if the measured average bodyweight is close to the target ($\pm 5\%$) and the flock is uniform.

In case the measured average bodyweight despite the ad libitum feeding is under the target weight, the feed intake has to be increased by running the feeders more often or by using higher nutrient feed formula until the target is reached.

In case the measured average bodyweight is above the target, the feed intake should not be increased until the target is reached. Never reduce the daily feed intake.

Grower feed can be used on the 8th week if the body weight of the pullets is around 700 g. Normally by the 15th week the average body weight of the pullets is around 1300 grams.

It is essential that TETRA-SL grows well from day one, as optimum skeletal development and adequate deposition of muscular and adipose tissues must be completed before the bird commences sexual maturation. It therefore follows that the shape of the body weight curve will depend on the anticipated age for

sexual maturity. The bird must develop its reproductive organs in the period 15-20 days before it lays its first egg, irrespective of when maturity occurs. The ovary and oviduct will weigh about 100-150 grams, and this will be in addition to the 8-12 grams daily growth that takes place at about this age. Body weight at first egg will therefore increase by 80-90 grams for each 10-day delay in maturity.

Once an individual bird starts to lay it is quite normal for its body weight to plateau, or even fall, for several weeks after first egg. In the period when a flock is starting to produce there will be three types of bird, (a) birds that have yet to develop sexually and growing at 8-10 grams per day, (b) birds undergoing development of the reproductive system and growing at 15 grams per day and (c) birds that are in lay, and having little or no body weight gain. The effect that this has upon the mean daily growth rate of a flock is shown in the graph describing the daily body weight gain before and after 50% rate of lay.

Body weight at first egg is a key factor in the determination of egg weight. The average egg weight in the laying year increases by 1.4-1.5 grams for each 100-gram increment of body weight.


It is a common phenomenon in brown egg laying birds that at about 15 weeks of age birds voluntarily reduce daily feed intake, and as a consequence reduce daily body weight gain. It follows therefore that body weights must be on target by this age if suboptimal weights are to be avoided. Birds, which commence egg production with body weights below optimum, frequently dip in egg production at or soon after peak rate of lay.

Oestrogen production by the ovary causes a retardation of skeletal growth. It is therefore extremely important that the skeleton has been allowed to develop optimally before the bird commences lay. Failure to permit satisfactory bone growth during the rearing period will predispose both skeletal disorders and shell quality problems during adult life.

The importance of obtaining the correct body weights for given ages and stages of sexual development dictates that sample body weighing of the flock is conducted at regular intervals during the rearing period. It is particularly important that the body weight is on target before a flock is changed from the Chick Starter to the Grower ration. It is also vital that early maturing flocks are fed ad libitum throughout the rearing period if the extra body weight gain necessary for these birds is to be achieved.


It will be seen from the TETRA-SL Body Weight Guide that early maturing flocks will initially have a faster body weight gain than later maturing flocks. Because individual body weights tend to plateau as soon as birds come into production and sexually immature birds continue to gain weight, the flock body weight averages cross over. As a result the earlier maturing birds then become lighter than later maturing birds. The body weight differences will persist throughout the life of the flock.

TETRA-SL Body Weight (Early, Standard and Late Maturity)



Pre-Lay (16 - 18 weeks)

In the 2-3 weeks before first egg enormous physiological changes take place within the bird. During this time the bird is depositing medullar bone, which will be involved in shell calcification once the bird is in lay. It is vital, therefore that



calcium intake is optimal at this stage if shell quality and bone strength is to be maximized later in the laying cycle. There is also an increased demand for energy and amino acids to furnish the development of the ovary and oviduct. Individual daily body weight gain will increase from about 10 grams per day to as much as 15 grams per day in the period 3-15 days before first egg. As the body weight has to be between 1500-1550 g before starting of lay, it is most important that feed intake is ad libitum whilst this ration is being fed, and that management factors, which depress feed intake are avoided or their consequences minimized as much as possible.

Because reproductive development and medullar bone deposition have a temporal relationship to sexual maturity, any change in flock maturity must be matched by a similar change in the introduction of the Pre-Lay ration.

Layer 1 (>19 weeks)

Early in the laying cycle feed intake is often relatively low, yet the demands for egg mass output and the continued increase in body weight will be at peak. Layer 1 ration must therefore be adequately formulated to satisfy the large demand for nutrients, when daily feed intake is probably less than 114 grams.

It is important that a layer ration includes a source of calcium, which is hard and relatively 'large', e.g. oyster shell, to provide calcium during the part of the day when the bird is not feeding. This will be in addition to a more soluble form of calcium carbonate, e.g. limestone flour, which will be more readily absorbed across the gut wall. Both forms of calcium are required in the ration to maximize shell quality.

Where it is not possible to feed a Pre-Lay ration Layer 1 may be introduced 3 weeks before anticipated start of egg production.

Layer 2 (>60 weeks)

In the later stages of the laying cycle egg mass output is decreasing, body weight gain is almost completed and feed intake is adequate. At this stage it is possible to minimize feed costs by feeding a lower nutrient density by controlling the ratio of the components.

TETRA-SL Feed Intake

Feed intake is greatly influenced by environmental factors and changes in consumption levels, without changes in formulation, will result in changes in egg output.

Laying hens primarily alter their daily feed intake to accommodate changes in their requirement for energy. Therefore factors such as ambient temperature, which alter the bird's demand for energy, automatically alter the bird's intake of feed. If there has been no modification of the ration formulation, changes in daily intake will result in changes in all nutrient intakes, and changes in the intake of amino acids, vitamins, minerals, yolk colorants, anticoccidials etc. will correspondingly affect bird performance.

The energy level of the ration itself will influence feed intake. TETRA-SL will reduce feed intake when the energy concentration of the ration is increased, and conversely increase intake when the energy level is decreased.

The laying hen does not totally adjust intake at the extremes of temperature or dietary energy. At high temperatures or with high-energy concentrations it tends to reduce energy intake too much and egg output suffers. At very low ambient temperatures, or on very low energy rations, the bird does not take in enough energy, and egg output is again reduced. Energy/amino acid ratios should be kept constant when the energy level of the ration is changed to ensure a satisfactory intake of protein, and amino acid concentrations should be increased when increasing temperatures result in lower daily feed intakes.

BODY WEIGHT GUIDE

Depth of feed in the trough, number of feeding per day and texture of feed affect feed intake. Consumption levels are positively correlated with depth of feed in the trough and feeding per day. Increasing depth of feed and number of times per day that feed is given to the birds will help maintain feed intake during hot weather. Birds also tend to eat more feed when it is given in a crumbed or pelleted form.

Boredom and copy feeding are factors, which tend to increase the feed intake of birds kept in cages. This is ironic, as caged birds require less energy for body temperature maintenance and activity than birds kept under extensive systems.

Feed intake is correlated with day length. TETRA-SL will consume about 1.5 grams more per day feed for each extra hour of light. This will not be totally wasted as egg numbers will be increased (about 3 eggs to 72 weeks) and average egg weight increased (about 0.1 grams per egg).

Feed consumption is also influenced by changes in sexual maturity. The combined effect of daylength (which is involved in determining age at first egg) and sexual maturity is shown in the following table.

Effect of age at first egg and daylength on daily feed intake (% change from 150 days and 12 hours)

Early maturing birds have smaller body weights during the laying year and produce less total egg output than later maturing birds, both factors reduce nutrient demand.

TETRA SL Body Weight Guide

Age (weeks)	Sexual maturity		
	Early (g)	Standard (g)	Late (g)
1	70	70	70
2	125	125	125
3	195	195	195
4	280	280	280
5	380	380	390
6	480	480	480
7	585	585	585
8	690	690	690
9	790	790	790
10	885	885	885
11	975	975	975
12	1 060	1 060	1 060
13	1 140	1 140	1 140
14	1 220	1 220	1 220
15	1 295	1 295	1 295
16	1 375	1 370	1 370
17	1 455	1 440	1 440
18	1 555	1 520	1 515
19	1 655	1 600	1 585
20	1 725	1 700	1 670
21	1 750	1 800	1 770
22	1 770	1 870	1 870
23	1 790	1 900	1 935
24	1 810	1 920	1 955
25	1 830	1 930	1 965
26	1 840	1 940	1 975
27	1 860	1 945	1 985
28	1 880	1 950	1 990
29	1 900	1 955	1 995
30	1 920	1 960	2 000

FEEDING PROGRAM (STANDARD MATURITY)

Age (weeks)	Body Weight (g)	Feed (g/bird/day)	Cummulative Feed (kg)	Type of Ration
1	70	9	0.06	Chick Starter
2	125	22	0.21	
3	195	26	0.40	
4	280	31	0.62	
5	380	36	0.87	
6	480	39	1.14	
7	585	44	1.45	
8	690	48	1.79	
9	790	51	2.15	Grower
10	885	54	2.53	
11	975	56	2.92	
12	1 060	59	3.33	
13	1 140	63	3.77	
14	1 220	68	4.25	
15	1 295	72	4.75	Pre-Lay
16	1 370	76	2.28	
17	1 440	78	5.83	
18	1 520	80	6.40	
19	1 610	Ad libitum feeding		Layer
20	1 725			

The feed quantities given in the table assume the use of rations, which have specifications similar to those detailed on page 15, feed intake and body weight gain resulting from it is affected by many nutritional and environmental factors (see Feed Intake section, page 10) As a result, feeding the quantities and types of rations described in the schedule will not necessarily produce the body weights desired. If body weights vary significantly from those given in the guide on page 11, appropriate changes should be made to the daily feed allocation.

If a non-standard maturity is chosen, the quantities of feed given above will need adjusting to produce the desired body weights. Early maturing birds will need more feed and late maturing birds will need less feed.

NUTRITION RECOMMENDATIONS

Ration	Chick	Grower	Pre-Lay	Layer 1	Layer 2
Age of bird (weeks)	0-8	9-15	16-18	19-60	>60
Metabolizable Energy (MJ/kg)	11.5	11.3	11.5	11.5	11.3
Metabolizable Energy (kcal/kg)	2 750	2 700	2 750	2 750	2 700
Crude Protein (%)	19	16	17	17.7	16.5
Amino acids					
Lysine (%)	1.00	0.65	0.80	0.85	0.75
Methionine (%)	0.45	0.35	0.35	0.40	0.35
Methionine + Cystine (%)	0.75	0.60	0.60	0.70	0.60
Tryptophan (%)	0.22	0.16	0.17	0.19	0.17
Threonine (%)	0.65	0.50	0.50	0.60	0.50
Minerals					
Calcium (%)	1.00	1.00	2.50	3.75	4.00
Phosphorus (total) (%)	0.70	0.65	0.70	0.70	0.65
Phosphorus (available) (%)	0.45	0.40	0.42	0.42	0.40
Sodium (%)	0.16	0.16	0.16	0.17	0.17
Fatty acids					
Linoleic (%)	1.25	1.00	1.20	1.30	1.30
Added Vitamins					
Vitamin A (iu/kg)	12 000	10 000	12 000	12 000	12 000
Vitamin D ₃ (iu/kg)	3 000	2 500	3 000	3 000	3 000
Vitamin E (iu/kg)	30	20	25	25	25
Vitamin K (mg/kg)	3	2	2	2	2
Thiamine (B ₁) (mg/kg)	2	2	2	2	2
Riboflavin(B ₂) (mg/kg)	6	4	6	6	6
Pyridoxine (B ₆) (mg/kg)	4	2	3	3	3
Vitamin B ₁₂ (mcg/kg)	20	10	20	20	20
Pantothenic Acid (mg/kg)	12	8	8	8	8
Niacin (mg/kg)	40	30	30	30	30
Biotin (mcg/kg)	100	100	100	100	100
Foline (mg/kg)	2	1	1	1	1
Choline (mg/kg)	400	300	400	400	400
Added Minerals					
Manganese (mg/kg)	100	100	100	100	100
Zinc (mg/kg)	80	80	80	80	80
Iron (mg/kg)	50	50	50	50	50
Copper (mg/kg)	8	8	8	8	8
Iodine (mg/kg)	1	1	1	1	1
Selenium (mg/kg)	0.30	0.30	0.30	0.30	0.30

LIGHTING PROGRAM

TETRA-SL will perform extremely well under a wide variety of lighting conditions. It is probably incorrect to talk about a specific lighting program for TETRA-SL, but rather to consider a lighting program as being specific for the production of a particular package of egg output. Program should therefore be designed to give the performance that best suits market demands.

The principle function of a lighting program is to influence the age at which a flock of birds becomes sexually mature. Age, and more particularly body weight, at first egg is the main factor, which determines the package of egg output. Egg numbers during the laying year decrease by 3-4 eggs for each 10-day delay in age at first egg. Average egg weight over the laying year will increase by about 1.4 g for each 10-day retardation of sexual maturity.

Standard lighting program

It is desirable, even when no specific change in maturity is required, that the lighting program includes a sufficiently large increase in daylength to trigger a uniform development of the ovary and reproductive tract within the flock. Small increases in daylength result in a wide range of individual ages at first egg, and this makes nutritional management of the flock more difficult. Larger light increases stimulate feed intake at a time when there is a massive increase in nutrient demand for growth of the reproductive organs, and for deposition of medullar bone. Lighting program is only effective if direct sunlight is blocked out of the building otherwise the time of maturity can vary. Due to this reason flocks moved to laying in autumn will start produce eggs a little later than stated in this management guide.

Lighting program for increasing egg weight

TETRA-SL will mature at about 19-20 weeks of age, even when no light increase is given. It is therefore

not possible to delay maturity by delaying the introduction of light increases. To successfully retard maturity in the TETRA-SL it is important that birds are reared on a decreasing light program during the first part of the growing period. The amount of delay in maturity depends on how long it takes to reach minimum daylength. Each extra week of decreasing daylength will result in a further one-day delay in age at first egg. It is important to note that sexual maturity delayed by controlling body weight gain will not result in any increase in egg weight.

Lighting program for increasing egg numbers

TETRA-SL can be readily brought into lay at an earlier age than normal by giving light increases at a younger age. However, it is really body weight at first egg, which controls egg output. Therefore it is vital that early maturing birds are fed ad libitum especially once a light increase has been given. It is not recommended to give increases in daylength before 15 weeks of age.

Initial light increase before maturity

It is possible to give the increase in daylength, which is given to initiate sexual maturity, as one abrupt jump rather than a series of weekly increases. This has the benefit of stimulating feed intake at a time when nutrient demand is increasing rapidly.

Maximum daylength

TETRA-SL will perform satisfactorily on a 12-hour light 12-hour dark light cycle. The provision of longer periods of light will increase both egg numbers and egg weight. The decision will be an economic one, as feed intake will increase by about 1.5 grams per day for each extra hour of light. Mortality rates are also likely to be higher, and recent findings suggest that increasing the light period may adversely affect shell quality. However, 16-17 hour daylengths make help maintain appetite during periods of high temperature.

LIGHTING PROGRAM FOR DIFFERENT MATURITY

LIGHTING PROGRAM FOR DIFFERENT MATURITY

Age	Sexual Maturity		
	Early	Standard	Late
0-2 days	23	23	23
3-5 days	14	14	18
6-8 days	12	12	18
9-11 days	10	10	16
12-14 days	8	8	16
3 weeks	8	8	15
4 weeks	8	8	14
5 weeks	8	8	13
6 weeks	8	8	12
7 weeks	8	8	11
8 weeks	8	8	10
9 weeks	8	8	9
10 weeks	8	8	8
11 weeks	8	8	8
12 weeks	8	8	8
13 weeks	8	8	8
14 weeks	8	8	8
15 weeks	8	8	8
16 weeks	11	8	8
17 weeks	12	8	8
18 weeks	13	9	8
19 weeks	14	10	9
20 weeks	14	11	10
21 weeks	14	12	11
22 weeks	14	13	12**
23 weeks	14	14	13
24 weeks onwards	14	14	14

***If egg weight becomes important after egg production has started daylength may be increased by 1 h per week to 16 hours.*

VETERINARY CONTROL

TETRA-SL hybrids can only perform their genetic potential if the risk of an infection of any kind of poultry disease is minimized. This risk can be minimized by keeping the following rules:

- Follow a thorough clean-out program before placing a new flock
 - Remove equipment and supplies from the building – wash, sanitize and store them in a clean area.
 - Remove all manure, old feed and rubbish.
 - Clean out the house blow or sweep down the dust, feather and other accumulated rubbish.
 - Completely wash the house with high-pressure water.
 - Disinfect the house perfectly.
 - Apply insect and vermin control.
 - Fumigate with Formaldehyde.
 - Use dry and mould-free litter.
- Minimize the entrance for unauthorised people and vehicles.
- Isolate the flock from other especially older flocks
- Avoid mixed-aged-flocks on the same farm.
- Keep wild birds, mice, rats and insects off the building.
- Keep every building clean on the farm
- Regular veterinary control is essential.

Recommendation for Vitamin Treatments

Age	Vitamin	Method
4-6. Day	A-, D ₃ -, E- and B-vitamin complex	Drinking Water
4. Week	A-, D ₃ -, E- and B-vitamin complex	Drinking Water
7. Week (after selection)	A-, D ₃ -, E- and B-vitamin complex	Drinking Water
12. Week	A-, D ₃ -, E- and B-vitamin complex	Drinking Water
19. Week (after housing)	A-, D ₃ -, E- and B-vitamin complex	Drinking Water
In every 4-6 weeks during laying period	A-, D ₃ -, E- and B-vitamin complex	Drinking Water

For different location a different vitamin treatment and vaccination program has to be designed considering the maternal immunities, local disease exposures, available vaccines and local veterinarian regulations.

EGG NUMBERS

TETRA-SL has been bred to produce an extremely high number of eggs. Egg production, however, is a character, which has relatively low heritability. This means that environmental factors such as health, nutrition, lighting and temperature have a large effect upon the number of eggs actually produced.

Egg output is influenced by both dietary energy and protein. Increasing the protein concentration of the ration, and ensuring that amino acids are correctly balanced, will minimize, but not prevent, reductions in rate of lay when high temperatures cause a drop in feed intake.

Egg output is the product of egg numbers and egg weight, and has a curvilinear relationship with nutrient intake, in particular the daily intake of protein. For each additional increment of protein the bird progressively produces more egg output, until an optimum protein intake is reached, at which point egg output plateaus. Conversely for each incremental reduction in protein intake the bird will produce progressively less egg output. Initially when the supply of protein is limiting TETRA-SL will respond by reducing egg numbers and egg weight equally, but when protein intake is more than 10% below optimum the major response will be a reduction in rate of lay. The response to changes in nutrient intake will be the same whatever the cause. Factors, which affect nutrient intake, include disease, nutrient concentrations in the ration, feed palatability, temperature, day length, light intensity and age at first egg.

TETRA-SL will reduce its daily intake of energy by 15-20 kJ (3.6-4.8 kcals) for each 1°C (1.8°F) rise in ambient temperature between 20 and 26°C (68 - 79°F). Above 26°C (79°F) the bird changes to a predominantly evaporative type of cooling and the rate of reduction in energy intake progressively

increases to around 50 kJ (12 kcals) per 1°C (1.8°F) rise. Each gram of egg requires 28 kJ (6.7 kcals) of energy, therefore, reductions in energy availability for egg production will result in lower rates of lay. Increasing the energy concentration of the ration will not increase daily energy intake; instead it will reduce daily feed intake and depress egg production by limiting further the bird's intake of protein.

The following chart shows the combined effects of age at first egg (50% rate of lay) and daylength in the laying period on number of eggs produced to 72 weeks of age.

Light (hours)	Age at 50% rate of lay (days)		
	140	144	148
8	299	295	291
10	305	301	297
12	311	307	303
14	317	313	309
16	323	319	315

TETRA-SL increases its egg numbers by about 3 eggs for each one hour increase in daylength. It must be remembered, however, that longer daylengths also result in larger feed intakes; daily feed intake will increase by approximately 1.5 grams for each extra hour of light per day. Advances in sexual maturity will increase egg numbers by about 3-4 eggs to 72 weeks of age for every 10-day younger age at first egg. Earlier sexual maturity, achieved through modifications of the lighting program, also increases daily feed intake and reduces average egg weight. It is, therefore, very important that a full analysis of feed costs and egg prices is performed before a decision is made regarding the lighting schedule.

RATE OF LAY (STANDARD MATURITY)

Age (weeks)	Egg Production (Hen-Housed)			Egg Production (Hen-Day)		
	Percent	Eggs per Hen	Cummulative	Percent	Eggs per Hen	Cummulative
18	0.0	0.00	0.0	0.0	0.00	0.0
19	7.0	0.49	0.5	7.0	0.49	0.5
20	25.0	1.75	2.2	25.0	1.75	2.2
21	50.9	3.56	5.8	51.0	3.57	5.8
22	74.8	5.23	11.0	75.0	5.25	11.1
23	88.7	6.21	17.2	89.0	6.23	17.3
24	92.6	6.48	23.7	93.0	6.51	23.8
25	93.5	6.54	30.3	94.0	6.58	30.4
26	94.4	6.61	36.9	95.0	6.65	37.0
27	95.3	6.67	43.5	96.0	6.72	43.8
28	94.7	6.63	50.2	95.5	6.69	50.4
29	94.1	6.59	56.8	95.0	6.65	57.1
30	93.6	6.55	63.3	94.5	6.62	63.7
31	93.5	6.54	69.9	94.5	6.62	70.3
32	93.4	6.54	76.4	94.5	6.62	76.9
33	92.8	6.50	82.9	94.0	6.58	83.5
34	92.7	6.49	89.4	94.0	6.58	90.1
35	92.1	6.45	95.8	93.5	6.55	96.6
36	92.1	6.45	102.3	93.5	6.55	103.2
37	91.5	6.40	108.7	93.0	6.51	109.7
38	90.9	6.36	115.0	92.5	6.48	116.2
39	90.3	6.32	121.4	92.0	6.44	122.6
40	89.8	6.29	127.6	91.5	6.41	129.0
41	89.2	6.24	133.9	91.0	6.37	135.4
42	88.6	6.20	140.1	90.5	6.34	141.7
43	88.0	6.16	146.2	90.0	6.30	148.0
44	87.5	6.12	152.4	89.5	6.27	154.3
45	86.9	6.08	158.5	89.0	6.23	160.5
46	86.3	6.04	164.5	88.5	6.20	166.7
47	85.8	6.01	170.5	88.0	6.16	172.9
48	85.2	5.96	176.5	87.5	6.13	179.0
49	84.6	5.92	182.4	87.0	6.09	185.1
50	84.1	5.89	188.3	86.5	6.06	191.1
51	83.7	5.86	194.1	86.2	6.03	197.2
52	83.3	5.83	200.0	85.9	6.01	203.2
53	83.0	5.81	205.8	85.6	5.99	209.2
54	82.6	5.78	211.5	85.3	5.97	215.1
55	82.1	5.75	217.3	85.0	5.95	221.1
56	81.8	5.73	223.0	84.7	5.93	227.0
57	81.4	5.70	228.7	84.4	5.91	232.9
58	81.0	5.67	234.4	84.1	5.89	238.8
59	80.7	5.65	240.0	83.8	5.87	244.7
60	80.3	5.62	245.7	83.5	5.85	250.5
61	79.9	5.59	251.3	83.2	5.82	256.4
62	79.6	5.57	256.8	82.9	5.80	262.2
63	79.2	5.54	262.4	82.6	5.78	267.9
64	78.8	5.52	267.9	82.3	5.76	273.7
65	78.4	5.49	273.4	82.0	5.74	279.4
66	78.1	5.47	278.8	81.7	5.72	285.2
67	77.7	5.44	284.3	81.4	5.70	290.9
68	77.4	5.42	289.7	81.1	5.68	296.5
69	77.0	5.39	295.1	80.8	5.66	302.2
70	76.6	5.36	300.4	80.5	5.64	307.8
71	76.2	5.33	305.8	80.2	5.61	313.4
72	75.9	5.31	311.1	79.9	5.59	319.0
73	75.6	5.29	316.4	79.6	5.57	324.6
74	75.1	5.26	321.6	79.3	5.55	330.2
75	74.8	5.24	326.9	79.0	5.53	335.7
76	74.5	5.22	332.1	78.7	5.51	341.2
77	74.2	5.19	337.3	78.4	5.49	346.7
78	73.8	5.16	342.5	78.1	5.47	352.1
79	73.4	5.14	347.6	77.8	5.45	357.6
80	73.1	5.12	352.7	77.5	5.43	363.0

EGG WEIGHT

As a result of many generations of selective breeding TETRA-SL possesses excellent egg weight characteristics. However, the demands of egg markets worldwide are both varied and changeable. This means that the modern laying hen must be adaptable and responsive to management manipulations designed to change it's primary egg weight characteristics. TETRA-SL is extremely versatile, and well able to respond to the various lighting and nutritional treatments used to change egg weight.

The single most influential factor, which can change egg weight, is age at sexual maturity; delays in maturity increase egg weight and advances in maturity reduce egg weight. However, it is important to note that egg weight is only affected when changes in sexual maturity are effected by alterations to the lighting program (see Lighting section, page 14). Maturity, which is retarded by controlling feed intake, will not improve egg weight. This is because it is the larger body weight, and not older age, of sexually delayed birds, which increases egg weight. Therefore control of nutrient intake during the rearing period cannot affect egg weight, because there is no increase in body weight at first egg. Average egg weight over the laying year is increased by about 1.4 grams for each 10 day delay in age at first egg, and correspondingly decreased by about 1.4 grams for each 10 day advance in maturity. When lighting programs are used to alter sexual maturity it is most important that the flock is fed ad libitum, because birds brought into lay at an early age require nutrients to develop the ovary and oviduct at a younger age, and birds held back, to increase egg weight, must start lay at a heavier body weight (see Body weight section, page 8). Often birds will take about two weeks to settle into production, and during this period they will lay eggs slightly smaller than birds already in lay - this is perfectly normal.

Once in lay the number of hours of light each day will also influence egg weight. There will be an increase of

about 0.1 grams in average egg weight for each extra one hour of light per day, but this will be linked to a 1.5 grams per day increase in feed intake.

The following chart shows the combined effects of age at first egg (50% rate of lay) and daylength in the laying period on average egg weight to 72 weeks of age.

Light (hours)	Age at 50% rate of lay (days)		
	140	144	148
8	61.4	62.8	64.3
10	61.6	63.0	64.4
12	61.8	63.3	64.7
14	62.1	63.5	64.9
16	62.4	63.8	65.2

Egg weight changes can be effected by modifications of the layer ration. Maximum daily egg output will require the bird to consume about 1000 mg total Lysine and 500 mg total Methionine per day, but such intakes of amino acids are unlikely to be economic, due to the law of diminishing returns. Practical intakes to obtain the most economic egg weight will be about 950 mg total Lysine and 450 mg total Methionine per day. Linoleic Acid intake is also important for maximizing egg weight, but again it is unlikely that the amounts necessary to give the best response biologically are economically justified. A minimum inclusion rate of 1.3% is recommended for satisfactory egg weight (see Nutrition section, page 8).

When egg prices and raw materials costs suggest a reduction in protein are economically desirable, reductions in egg numbers are likely to be greater than reductions in egg weight, especially when the resulting amino acid intake falls below 90% of that necessary for maximum egg output.

EGG WEIGHT AND SIZES (STANDARD MATURITY)

Age (weeks)	Average Egg Weight (g)	Egg Sizes (% of all Eggs)			
		XL Extra Large >73g	L Large 63-73g	M Medium 53 - 63 g	S Small <53 g
19	45.2	0.0	0.0	5.7	94.3
20	48.1	0.0	0.0	13.7	86.3
21	50.4	0.0	0.9	28.6	70.5
22	52.7	0.0	1.5	45.1	53.4
23	55.0	0.0	7.3	55.9	36.8
24	57.3	0.1	12.2	67.5	20.2
25	58.8	0.3	21.7	62.6	15.4
26	60.2	0.8	29.1	61.5	8.6
27	60.9	1.3	34.3	58.3	6.1
28	61.6	1.5	38.0	55.2	5.3
29	62.0	1.6	40.6	53.7	4.1
30	62.4	1.9	42.3	52.1	3.7
31	62.7	2.0	45.6	49.9	2.5
32	63.0	2.2	48.1	47.5	2.2
33	63.3	2.4	50.0	45.7	1.9
34	63.6	2.7	52.7	43.0	1.6
35	63.8	2.9	53.2	42.5	1.4
36	64.0	3.0	55.9	39.8	1.3
37	64.2	3.3	57.1	38.5	1.1
38	64.4	3.5	58.4	37.1	1.0
40	64.8	3.9	60.6	34.5	1.0
42	65.2	4.7	62.1	32.3	0.9
44	65.5	5.5	63.9	29.7	0.9
46	65.8	6.1	64.2	28.9	0.8
48	66.1	7.4	64.5	27.4	0.7
52	66.5	8.6	64.6	26.2	0.6
56	66.9	9.3	64.8	25.4	0.5
60	67.1	10.7	64.9	24.0	0.4
64	67.3	11.8	65.0	22.9	0.3
68	67.4	12.6	64.8	22.3	0.3
72	67.5	13.4	64.7	21.6	0.3
76	67.6	13.9	64.5	21.3	0.3
80	67.7	14.2	64.2	21.3	0.3

EGG QUALITY

TETRA-SL has undergone rigorous selection for improvement in all internal and egg shell quality traits. However, egg quality is very complex and much research is still required to fully understand the biological mechanisms involved to egg formation. Many factors are known to affect shell quality in addition to the genetic make-up of the bird; age of bird, position of egg within the sequence, egg weight, bird behaviour, lighting regime, nutrition, disease, drugs, temperature, housing/egg collection system and feeding patterns.

The egg shell is not a uniform layer of calcium carbonate, but a complex multilayer structure. Defects on the shell surface can be the result of a poorly formed mamillary layer, this is the layer of cones which form the foundation of the shell structure, and/or an unsatisfactorily constructed palisade layer.

Calcium is the main mineral involved in shell deposition, but other nutrients play part too. These include Chlorine, Sodium, Potassium, Phosphorus, Vitamin D₃, Manganese, Magnesium, Fluorine and Zeolite A (a sodium alumino silicate compound). It is important to have the right balance of minerals as well as the correct absolute quantities. The Calcium:Phosphorus ratio, for example, needs to be widened as the flock ages, however, if too wide there may be adverse effects on rate of lay. Calcium particle size affects transit time in the gut, and absorption into the blood stream. Ideally at least half of the calcium carbonate in the ration should have a large particle size.


In the 15-20 days before an individual hen lays its first egg medullar bone will be deposited in the long bones of the skeleton. Medullar bone comes partly from the ration and partly from trabecular bone, and is the source of calcium for shell formation. It is vital therefore that the ration fed to the birds during the period approaching sexual maturity that calcium levels in the ration are increased (see Nutrition section page 8).

Poor shell quality may be improved by modifications of the lighting regime. Ephemeral lighting programs (e.g. 28 hour light-dark cycles), and repeating short cycles (e.g. a repeating 3-hour light, 3-hour darkness cycle) will increase shell weight and thickness. It is possible to interchange between these types of non-24 hour lighting and conventional lighting without any problems. However, when ephemeral and short cycle repeating patterns are used in the first half of the laying period there will be a reduction in rate of lay, but a compensatory increase in egg weight.

Recent research has indicated that long daylengths may be detrimental to shell quality. Shell weight appears to decrease with increasing daylengths, and when combined with the larger egg weight observed under long photoperiods, reduces shell thickness. Long photoperiods also increase the incidence of body-checked eggs. These are caused by contractions of the uterus (shell gland), when the shell of the egg is fragile, cracking the shell round the equator. Calcification then takes place on top of the cracks to form the typical equatorial bulge. Manipulations of the lighting program to minimize activity at the end of the light period will help alleviate this problem.

Shell quality will be reduced by high ambient temperatures, especially over 32°C (90°F) and high relative humidity. Any attempt to maximize nutrient intake will help to minimize the decline in quality; however, the problem is not only caused by low calcium intakes. A reduction in blood flow to the uterus caused by vasodilatation and a decrease in bicarbonate availability due to respiratory alkalosis will aggravate the problem.

EGG PRODUCTION AND EGG WEIGHT GRAPH



HOT CLIMATE MANAGEMENT

In the open house system of poultry keeping practised in tropical climates it is not possible for the TETRA-SL to express its genetic potential fully. However, there are various modifications to the management recommendations for controlled environment conditions, which will minimize the loss of performance.

The two main problems encountered when keeping birds at high temperatures are the difficulties of getting enough nutrients into the bird and the change in method of heat loss used by the bird to maintain its body temperature after panting begins at about 28°C (82°F). Laying hens reduce their intake of energy as ambient temperatures rise because of the lower demand for heat production at higher temperatures. The difficulty for the egg producer is that voluntary energy intake falls curvilinearly whilst heat production falls linearly, resulting in less energy being available for egg production at high temperatures, particularly above 28°C (82°F).

°C/°F	ME intake (kJ/d)	Heat loss (kJ/d)	Egg energy (kJ/d)
20/68	1260	953	307
22/72	1232	915	317
24/75	1198	876	322
26/79	1156	838	318
28/82	1105	800	305
30/86	1043	761	282
32/90	968	723	245
34/93	879	685	194
36/97	774	647	162
38/100	651	608	43

The table clearly illustrates the peak availability of energy for egg production at 24°C (75°F), and the decline in energy available for egg production after about 28°C (82°F).

There are two approaches to minimizing performance losses, (a) reducing the temperature of the bird's microclimate, and (b) maximizing nutrient intake. The first method ideally requires consideration before the poultry unit is built, as house design is involved. However, existing units can be modified to reduce the

effects of solar radiation. Obviously with open housing little can be done to reduce air temperature within the house. The second approach is probably the easier to implement, as alterations to ration specification and lighting schedules are already part of normal stock management.

Housing


1. Use roofing materials, which have good insulation properties and reflect solar radiation. Natural materials like palm thatch usefully reduce penetration of solar heat. Where metal roofs are unavoidable they can be painted with a solar reflective product, or be fitted with a sprinkler along the ridge to reduce the heat radiated through to the house interior. Where water is at hand premium sprinkler systems can incorporate a recycling pump.

2. Roof overhangs should be long enough to prevent direct sunlight falling on the birds, at least in the hottest part of the day. Ridge outlets should be fitted to permit bird heat to escape, ideally these should be open away from the sun. Roofs should be as high as possible to minimize the temperature at bird level and maximize the natural airflow to the ridge.

3. Mount fans vertically to create horizontal air movement at bird's level (Safety guards must be fitted). At air speeds of 2m/sec direct heat is doubled (plumage will be ruffled at this speed). This type of cooling must not be used when ambient temperature exceed 40°C (104°F).

4. Fogging the inside of the house with fine water droplets will reduce the air temperature and moisten the fleshy parts of the bird's head, so improving heat loss. Nozzle size should be small enough to ensure that the water droplet size is not too big; nozzles working at 140 kPa (20 lbs/in) will use 3.4 litres (0.75 gallons) per hour and produce a smoke-like mist.

5. Vegetation and trees may be planted around the buildings to provide shade and reduce the amount of



sunlight reflected from the ground. Care must be taken not to restrict the natural airflow into the house.

Bird management

1. Radiative heat transfer between birds will be reduced by providing the birds with more space. Stocking rates should be reduced by 2% for each 1°C (2°F) rise in temperature above 20°C (68°F).

2. Natural daylengths in tropical areas vary seasonally from 11-13 hours. An extension of the daylength into the cooler part of the day with artificial lighting will have two beneficial effects upon feed consumption. Firstly, feed intake is positively related to daylength, and secondly it is negatively correlated with temperature. It is, however, undesirable to have daylengths longer than 17 or 18 hours. If the provision of extra light is started before the birds start egg production, the increase in daylength will act as a stimulator of sexual maturity. The timing of this light increase should be carefully considered as early maturity will reduce egg weight, and reduced egg weight will already be anticipated because of high temperatures and low energy intakes.

Nutrition

1. Use any management technique, which stimulates feed intake; for example frequent feed deliveries. Avoid stale or unpalatable feed.

2. One of the major nutrients for consideration is water. Restrictions on water intake, whether caused by physical means, insufficient drinkers, high water temperature or poor quality, will adversely affect feed intake. This will obviously result in a reduction in all other nutrients. Water requirements increase curvilinearly with temperature (see table for an adult layer).

Air temp °C/°F	Water intake (ml/bird/day)	Water to energy ratio (ml/MJ)
20/68	210	165
25/77	250	210
30/86	320	305
38/95	420	505
40/104	535	1050

Obviously any factor, which restricts water intake, should be avoided. Water-cooling may be considered if feasible.

3. While temperatures are below 28°C (82°F) the concentration of metabolizable energy in the feed may be increased in proportion to the decline in feed intake. Beyond 28°C (82°F) energy levels in the ration should be reduced to encourage feed intake and facilitate inclusion of protein.

4. Protein requirements for egg production may be regarded as being independent of temperature. Rations should therefore be formulated to provide the amino acid intake required for the egg output potential made possible by the energy intake/heat production surplus (1 gram of egg requires 6.69 kJ energy).

BÁBOLNA TETRA Ltd.

H-9651 Uraiújfalu; Petőfi u. 18, Hungary
Phone: +36 95 345 002; +36 95 345 004

Fax: +36 95 345 009

e-mail: tetrakft@babolnatetra.com

www.babolnatetra.com